

## Methods of Detection Utilizing Modified Bacteriophage

### BACKGROUND OF THE INVENTION

5

#### Field of the Invention

The present invention provides improved assays for detection of proteins utilizing virus expressing a ligand for the protein. Such assays can be used in areas  
10 wherein traditional immunoassays have previously been utilized.

#### Background Art

Conventional immunoassays rely on the high affinity specific interaction between immunoglobulin and its corresponding epitope. Generally, a variable region of  
15 immunoglobulin forms a binding pocket that recognizes either conformation or primary sequence of a given epitope. Because the epitope can be a very small and linear region of a protein, it is common to find that an immunoglobulin recognizes a protein in either native or denatured conformation by binding to a small embedded peptide. Therefore detection agents that can recognize native conformations of peptides, such as receptors,  
20 channels, other surface proteins, and extracellular proteins, would be highly useful.

Use of random peptide phage libraries to isolate lead compounds for target proteins has yielded a variety of peptides that recognize specific target proteins with interesting biological activities. Such strategy has provided interesting binding peptides  
25 which recognize important proteins including antibodies (Smith, 1985; Cwirla et al., 1990; Scott and Smith, 1990; Kay et al., 1993), streptavidin (Devlin et al., 1990), calmodulin (Dedman et al., 1993), SH3 domain (Sparks et al., 1994), and endoplasmic reticulum protein BiP (Blond-Elguindi et al., 1993). Some of these peptides have yielded important insights into the structure-function of the target protein. For example,  
30 analysis of peptides that bind to BiP have shown that BiP preferentially binds peptides containing a subset of aromatic and hydrophobic amino acids in alternating positions,

suggesting that peptides bind in an extended conformation, with the side chains of alternating residues pointing into a cleft on the BiP molecule (Blond-Elguindi et al., 1993).

- 5 N-methyl D-aspartate receptor (NMDAR) channels, found in brain, belong to the superfamily of ligand-gated ion channels, which become selectively permeable to ions upon binding to ligands. The glutamate-mediated channel activity is important for synaptic plasticity, synaptogenesis, and excitotoxicity. NMDA receptors are encoded by at least five genes (NR1, 2A, 2B, 2C, and 2D) and their spliced variants <sup>7</sup>. NR1 is an
- 10 essential subunit for mediating glutamate-induced channel activity, since a homomultimeric NR1 complex, when expressed in *Xenopus* oocytes, exhibits some of key properties found in native channels. These properties include direct permeability of calcium, voltage dependent  $Mg^{2+}$  blockade of the ion channel, and binding sites for modulators such as  $Zn^{2+}$ , glycine, and polyamines. The other subunits by themselves
- 15 cannot form a functional channel. However, they coassemble with NR1 to form heteromultimers which increase the expression level and channel diversity. The NR1 polypeptide can be divided into two regions: an extracellular amino-terminal domain containing the putative agonist binding site, and a hydrophobic core region with multiple transmembrane segments which participate the formation of the ion conducting pathway.
- 20 Although several classes of compounds have been found to modulate the NMDA receptor, at the molecular level little is known in terms of amino acid residues that are responsible for the interaction. To better understand the physiological roles of NMDA receptors with different subunit composition or to detect NMDA receptors in samples, it would be helpful to develop specific ligands that are capable of distinguishing different
- 25 subunits.

Excitotoxic mechanisms have a well established role in the pathogenesis of neuronal injury following acute CNS insults such as ischaemia and trauma. Increasing evidence now supports their roles in chronic neurodegenerative disorders such as motor

30 neuron diseases (MND). Although the molecular mechanisms underlying excitotoxic neuronal injury are still being elucidated, a large body of evidence indicates that the

cascade of events resulting from elevation of intracellular free calcium is likely to play a major role, which is thought to be mediated by NMDA receptor and voltage-gated calcium channels. NMDA receptor has several very important properties of relevance to excitotoxicity, particularly, the ability to permeate calcium ions. Thus, understanding of

5 NMDA receptor properties at the molecular level and development of reagents that specifically modulate these receptors can be of importance for understanding excitotoxicity resulted diseases and providing potential leads which could facilitate future treatments. By further improving the affinity and stability of NMDA receptor-specific ligands, their potential neuronal protective activity can be improved.

10

The present invention provides new procedures which use bacterial virus (including bacteriophage) as a detective reagent, in a way comparable to primary antibody in immunoassays, to monitor the expression of receptor and channels in mammalian cells. Conversely, the phage binding described herein requires the

15 corresponding receptor in proper conformation to present the binding pocket for the peptide to bind. Because the conformation of receptor binding pocket is critical for the peptide binding, peptide-receptor target interaction per se can have higher specificity than that of some antibody-receptor interaction. In search of novel peptides that modulate receptor activity, phage clones with specific interacting peptides from random

20 peptide libraries have been isolated by panning selection, which is often based on the multivalent interaction between the phage particle and target receptor. These assays provide an improved way to detect protein expression and to study peptide-receptor interaction at the molecular level.

25

## BRIEF DESCRIPTION OF THE DRAWINGS

- Figure 1. Isolation of Mag-4 peptides that specifically associate with N-NR1. (A) Amino acid sequence alignment. Two phage clones, Mag-4.1 and Mag-4.2, specifically association with N-NR1 were isolated by panning selection (see Experimental Protocols). The amino acid sequence was deduced and aligned. (B) Binding specificity of Mag-4.1 peptides to the NH<sub>2</sub>-terminal domain of glutamate receptor. Phage ELISA experiments were carried out using N-NR1, N-NR1.spl, and N-GluR1. Experimental procedures were described in Experimental Protocols. The dash bars indicate wells coated with bovine serum albumin (BSA) only, white bars: BSA plus mAb8610; black bars: BSA, mAb8610 plus the captured fusion proteins for N-NR1, N-NR1.spl, or N-GluR1 (as indicated). Experiments were carried out in triplicate. Error bars indicate the SD (C) Inhibition of Mag-4.1 phage binding by synthetic Mag-4.1 peptide. The ability of synthetic HPLC-purified Mag-4.1 peptide (GGGLNDWFITYIGGG) (SEQ ID NO:4) to compete with the Mag-4.1 phage binding to N-NR1 fusion protein was determined by phage ELISA. The vertical axis indicates the relative binding comparing to the signal obtained without adding the peptide inhibitors. The horizontal axis indicates concentration of the peptides added in mM (as indicated).
- Figure 2. Specific detection of N-NR1 protein by phage-overlay. Crude extracts containing fusion proteins of N-NR1 and N-NR2A at 1:5 ratio were fractionated on SDS-PAGE and transferred onto a nitrocellulose filter. The immobilized proteins were subjected to a denaturation-renaturation procedures (see Experimental Protocol). The renatured protein were detected by either monoclonal antibody 8610 (mAb8610) (left panel) or Mag-4.1 phage (right panel). Lanes 1, 2, 3 and 4 contain decreasing amounts of N-NR1 fusion protein (100 ng, 20 ng, 4 ng, and 0.8 ng).

- Figure 3. Phage-immunostain detection of N-NR1 protein on cell surface. Two CHO cell lines expressing low ( $\sim 10^4$  copies per cell) and high ( $\sim 10^6$  copies per cells) of N-NR1 fusion protein were mixed by 1:1 ratio and allowed to grow on gelatin-coated cover glass. The mixed cells were incubated with Mag-4.1 phage in the presence (A) or

absence (B) of 500 mM of synthetic Mag-4.1 peptide. The Mag-4.1 phage binding was detected by rabbit-anti-phage antibody (see Experimental Protocol).

Figure 4. Double stain of N-NR1 protein by mAb8610 and Mag-4.1 phage. N-NR1 positive and N-NR1 negative cells were mixed at ratio 1:1 and allowed to grow on gelatin-coated cover glass. The live cells were first incubated with Mag-4.1 phage. After fixation, the cells were incubated with mAb8610 (mouse) specific to the epitope on fusion protein and anti-phage antibody (rabbit). The antibody binding was then detected by goat-anti-mouse (Texas-red) and goat-anti-rabbit (fluorescein, green). A same area of cells were visualized by (A) Nomaski, (B) mAb8610; and (C) Mag-4.1.

Figure 5. Detection of the N-NR1 protein by FACS. CHO cells were stained with Mag-4.1 phage followed by rabbit anti-phage antibody and fluorescein-labeled secondary antibody (see Figure 4 legend and Experimental Protocol). Vertical axis is relative cell number, horizontal axis is relative intensity of green fluorescence. (A) CHO cells ; (B) CHO cells expressing low level of N-NR1 protein; (C) same as B, but stained in the presence of 500 mM of Mag-4.1 peptide; (D) a 1:1 mixture of CHO cells expressing high and low levels of N-NR1 protein; (E) same as D, but in the presence 500 mM of Mag-4.1 peptide.

Figure 6. Detection of functional NMDA receptor by Mag-4.1 phage. HEK cells were cotransfected with plasmids expressing NR1 and NR2A cDNAs to form functional NMDA receptor channels (Li et al., 1996). At 48 hours after transfection, the cells were subjected to phage-immunostain using Mag-4.1. The phage binding was visualized by HPR-cojugated antibody (see Experimental Protocol). Same area of stained cells was shown in low magnification (top panel) or high magnification (lower panel).

## DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a method of detecting the presence of a polypeptide in a sample comprising contacting with the sample a detectable virus  
 5 expressing on its surface a ligand for the polypeptide and detecting binding of the virus to the sample, thus detecting the presence of the polypeptide in the sample.

By "sample" is meant a sample wherein the selected polypeptide in the sample is not an isolated polypeptide, some examples of such a sample including a protein  
 10 homogenate from a cell, tissue or animal; cells; tissue section; biological or clinical sample such as blood, urine, saliva; any other sample to be tested for presence of contaminants comprising proteins (e.g., water sample, food sample). For example, the presence of a microbe can be detected in a food, water or clinical sample by detecting the presence of a protein, intracellular, surface or extracellular, that is a component of  
 15 the microbe in the sample.

As used in the specification and in the claims, "a" can mean one or more, depending upon the context in which it is used.

20 The terms "polypeptide" and "peptide" refer to a polymer of amino acids and includes full-length proteins and fragments thereof.

By "selected" polypeptide is meant that the target protein one is directing the phage or virus to bind is known. For example, a selected target can be the NMDA  
 25 receptor, and therefore a ligand that can be used to detect the presence of the receptor in a sample can be, e.g., Mag-4.1 or Mag-4.2, which are shown herein to be ligands for NMDA receptor. Thus a selected polypeptide is predetermined prior to detecting the presence of the polypeptide by detecting binding of the phage or virus expressing the ligand to the polypeptide.

30

The virus utilized in the method can be a bacteriophage. For example the bacteriophage can be bacteriophage f1, M13, and other bacteriophages known in the art. Viruses can include any other desired virus, as will be recognized by those of skill in the art, such as adenovirus, etc. The phage or virus can be modified in any of various ways  
 5 known in the art, such as to be rendered replication-deficient or to eliminate other viral genes, and methods of such modifications are standard in the art. The virus is modified to express the ligand on the surface of the virus, as by engineering the virus genome to encode a fusion protein for a coat protein and the ligand. For example, in bacteriophage, the ligand can be encoded by pIII or pVIII protein. Making  
 10 bacteriophage that express peptides on their surface is standard in the art (described in *e.g.*, US Pat No. 5427908; US Pat No. 5403484; US Pat No. 5432018; US Pat No. 5432018; US Pat No. 5270170; WO 92/06176; Smith et al. (1993); Kolvunen, E. et al., *J. Cell Biol.* 124:373 (1994); Kolvunen, E. et al., *Meth. Enzymol.* 245:346 (1994); Pasqualini, R. et al. *J. Cell Biol.* 130:1189 (1995)).

15

The present invention includes that the protein can be a cellular protein. Cellular proteins can include intracellular proteins, extracellular proteins, and/or surface proteins. Cellular proteins as used in these methods can also include viral proteins expressed by a cell that has been infected by a virus.

20

The sample in which the presence of the polypeptide is detected can be a clinical sample. Clinical samples can include any relevant sample from the body, such as excised tissue, blood, plasma, urine and saliva, and protein extracts from any such sample. Other samples can include cultured cells, proteins secreted by cells, and the like.

25

Specific binding between an receptor or channel and a ligand means that the ligand can be used to selectively remove the receptor or channel from a sample or to inhibit the receptor or channel's function and can readily be determined by radio immune assay (RA), bioassay, or enzyme-linked immunosorbant (ELISA) technology using an  
 30 antibody, preferably monoclonal, specific for the ligand.

The present invention provides a method of detecting the presence of a selected polypeptide in a sample comprising contacting with the sample a detectable virus, including a bacteriophage, expressing on its surface a ligand previously demonstrated to specifically bind the selected polypeptide and detecting binding of the virus to the sample, thus detecting the presence of the selected polypeptide in the sample.

A ligand can be previously determined to specifically bind the selected protein by any known, standard means for determining such binding or, for example, as described herein. A ligand can include, for example, a peptide hormone, a toxin, a fragment from a large protein.

A detectable virus or phage is one that can be detected by any of many possible means. For example, the phage or virus can be detected by a polyclonal or monoclonal antibody directed against the virus by the antibody itself being labeled or by utilizing a secondary antibody that is labeled or has a recognition site for a label. Additionally, the phage itself can be directly detectable, as described herein.

The present invention further provides a method of detecting the presence of a selected cellular protein on the surface of a cell comprising contacting with the cell a detectable virus expressing on its surface a ligand previously demonstrated to specifically bind the selected cellular protein and detecting binding of the virus to the cell, thus detecting the presence of the selected cellular protein on the surface of the cell. Such detection includes that the cell can be *in vitro* or *in vivo*.

Conditions for contacting phage or virus with the selected polypeptide can be selected based upon the location of the target. In general, conditions for phage-receptor (or other cellular protein) binding can be as described in Sparks, A.B., *et al.* (1995) *Methods. Enzymol.* 255:498-509. For example, *in vitro* conditions for binding standard detection reagents, such as monoclonal antibodies, to cells, or for phage-receptor binding, for binding reagents to a protein blot (*e.g.*, Li, *et al.* (1992)), for ELISA (*e.g.*, Barrett, *et al.* (1992)), for binding a reagent to a tissue sample, can be



followed, with any modifications desired, particularly in light of typical conditions for binding phage libraries and in light of the examples provided herein. Additionally, *in vivo* conditions for contact can be determined by conditions utilized for binding, e.g., monoclonal antibodies and viruses *in vivo*. Virus can be administered via the

5 bloodstream, as by intravenous administration or by administering to an artery supplying a target organ, or by direct injection into a target organ, by inhalation or intratracheal injection for delivery to the lungs, and conditions can be modified accordingly such as for administration of viruses *in vivo* for, e.g., gene therapy.

10 The present invention provides a method of detecting the presence of a selected polypeptide in a sample comprising contacting with the sample a detectable bacteriophage expressing on its surface at least 10, or at least 100, or at least 400, or at least 800, or at least 1000 copies of a ligand for the selected polypeptide and detecting

15 binding of the bacteriophage to the sample, thus detecting the presence of the selected polypeptide in the sample. Typically, by encoding the ligand within the pVIII coat protein, the ligand is expressed in high copy number. The present invention further provides a method of detecting the presence of a selected cellular protein on the surface of a cell comprising contacting with the cell a detectable bacteriophage expressing on its

20 surface at least 10 copies, or at least 100, or at least 400, or at least 800, or at least 1000 copies of a ligand for the selected cellular protein and detecting binding of the bacteriophage to the cell, thus detecting the presence of the selected cellular protein on the surface of the cell.

The present invention further provides a method of isolating a cell expressing a

25 selected polypeptide on its surface comprising contacting with a sample of cells a virus expressing on its surface a ligand previously demonstrated to be specific for the selected polypeptide and isolating bound virus, thereby isolating a cell expressing the selected polypeptide on its surface. The cell that binds the virus is isolated, for example, by a fluorescence activated cell sorter if the virus is labeled, either directly or indirectly, by a

30 fluorescent moiety. For example, a cell expressing the NMDA receptor can be isolated by utilizing a virus or bacteriophage expressing on its surface Mag-4.1 or Mag-4.2.

The present invention further provides a virus, including a bacteriophage, comprising a label such that the label can be directly detected. The virus can express on its surface a ligand for a selected protein. By "directly detected" is meant that the bacteriophage can be labeled in advance and still bind its target upon addition to a cell/sample, etc. containing the target for direct detection via the label on the bacteriophage, *i.e.*, no secondary antibody is necessary. Furthermore, by "label" is meant a means for visualization, such as a recognition site for direct phosphorylation, biotinylation, chemical linkages, etc. engineered into the virus, or such as a directly visualized label requiring no chemical reaction to detect, *e.g.*, the virus expresses a fluorescent protein or is labeled by a radioactive moiety. Phage can be modified to include a label in advance, then bound to the phage target in a sample, and visualized to detect the presence and localization of the bound phage. Visualization may or may not require a chemical reaction. Thus, directly detected does not require interaction with a secondary labelled ligand, for example an antibody. An example of a fluorescent protein is the green fluorescent protein (GFP) originally isolated from the jellyfish *Aequorea victoria*. Another example of a fluorescent protein is the green fluorescent protein originally isolated from *Renilla reniformis*, which demonstrated a single absorption peak at 498 nm and an emission peak at 509 nm. (Cubitt, *et al.* (1995) TIBS 20: 448-455). Phage expressing, for example, a peptide hormone, a toxin, or a fragment from a large protein can, for example, be used for a drug screening procedure.

A peptide, including a bacteriophage expressing the peptide, can be administered to a subject, including a human or an animal model, by any of many standard means for administering proteins to cells and to animal subjects. Typically, peptides can be administered orally, parenterally (*e.g.*, intravenously), by intramuscular injection, by intraperitoneal injection, topically, transdermally, or the like. Peptides can be administered, *e.g.*, expressed on the surface of a bacteriophage (such as bacteriophage f1 or M13), expressed on the surface of a cell, or incorporated into a lipid composition such as a cationic liposome.

Virus or phage compositions can include various amounts of the selected virus in combination with a pharmaceutically acceptable carrier and, in addition, if desired, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, diluents, etc. Parental administration, if used, is generally characterized by injection. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms  
 5 suitable for solution or suspension in liquid prior to injection, or as emulsions. (as described in *e.g.*, Martin, E.W. (ed.) *Remington's Pharmaceutical Sciences*, latest edition Mack Publishing Co., Easton, PA.).

10

To obtain a peptide ligand for a given protein target, a typical experimental paradigm involves three steps. First, phage clones carrying lead sequences are isolated from random peptide virus or phage libraries based on specific binding to the purified target protein. Second, according to the amino acid sequence information from the lead  
 15 peptide sequences, mutagenised libraries can be constructed to identify homologous sequences with higher affinity. Third, based on the amino acid sequence, synthetic peptides can be obtained to directly test their affinity and activity in bioassays. Since the region or surface present in the putative functional domain may not be accessible in native receptor, it can be particularly useful to be able to evaluate binding activity of  
 20 peptides present on candidate phage clones during early stage of phage selection.

As will be recognized by those skilled in the art, numerous types of immunoassays are available for use in the present invention to detect binding between a protein, such as a receptor or channel, and a peptide or other ligand of this invention.  
 25 For example, a monoclonal antibody specific for the phage bearing the peptide can be contacted with the bound sample followed by secondary antibody and standard imaging. Direct and indirect binding assays, competitive assays, sandwich assays, and the like, as are generally described in, *e.g.*, U.S. Pat. Nos. 4,642,285; 4,376,110; 4,016,043; 3,879,262; 3,852,157; 3,850,752; 3,839,153; 3,791,932; and Harlow and Lane, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, N.Y. (1988).  
 30 Methods of producing antibodies to detect any phage utilized herein are standard in the

art (*see. e.g.* Harlow and Lane, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, N.Y. (1988).; WO 93/06213).

The examples herein show the isolation of a phage clone carrying a low affinity  
5 peptide, which specifically recognizes truncated NMDA receptor fragment. By taking  
advantage of the specific binding between the purified phage particles and receptor, the  
present invention provides phage-immunoblot, phage-immunostain, and phage-FACS  
procedures to monitor expression of NMDA receptor protein, or any other selected  
protein (with corresponding ligands on the phage), in mammalian cells. These  
10 procedures offer a new way to help monitoring receptor protein expression. In addition  
they can facilitate molecular characterization of the specific peptide-receptor interaction.

In order to detect an intracellular target, some modifications can be made to  
improve the penetration of phage particle to reach target proteins. Such improvement  
15 can involve, for example, partial disruption of phage filament to generate shorter  
truncated phage filaments which have better penetration but retain multivalent  
peptide-receptor binding. Furthermore, if the specificity of phage staining is affected by  
the potential interaction between phage coat proteins and other cellular proteins, a phage  
variant with altered coat proteins that have lower background binding can readily be  
20 developed given the teachings herein. Phage variants can readily be tested for such  
reduced background following the methods herein.

#### Statement Concerning Utility

Monitoring receptor expression using recombinant bacteriophage with specific  
25 peptide ligands shares several advantages of other immunoassays. In particular, the  
present assays take advantage of the existing signal amplification strategies, since the  
binding of phage can, if desired, be eventually visualized by enzyme- or fluorescent dye-  
conjugated secondary antibodies that are available commercially. It is important,  
however, to note that there is fundamental difference between the phage-mediated  
30 immunoassay and the conventional primary antibody-mediated immunoassays.  
Generally, a variable region of immunoglobulin forms a binding pocket that recognizes

either conformation or primary sequence of a given epitope. Because the epitope can be a very small and linear region of a protein, it is common to find that an immunoglobulin recognizes a protein in either native or denatured conformation by binding to a small embedded peptide. Conversely, the phage binding described herein requires the

5 corresponding receptor in proper conformation to present the binding pocket for the peptide to bind. Because the conformation of receptor binding pocket is critical for the peptide binding, peptide-receptor target interaction per se can have higher specificity than that of some antibody-receptor interaction.

10 For peptides with interesting biological and/or therapeutic activity, it is important to understand their action in the molecular and mechanistic detail. One of the first steps to address this question is to determine the critical interacting residues on both receptor and peptides. Use of bacteriophage as a vehicle to carry multiple copies of receptor

15 interacting peptide offers several interesting approaches for detailed molecular analysis of peptide-receptor interaction, which should be applicable to both an existing bioactive peptide (e.g., angiotensin II) or novel lead peptides (from random peptide libraries). A combination of phage-FACS analysis and random mutagenesis strategy (e.g., known strategies such as described in Rice et al., 1992) on the target receptor can offer a

20 powerful approach to select interacting mutated receptors, which can lead to determination of interacting residues that mediate the phage-receptor interaction.

The present invention is more particularly described in the following examples which are intended as illustrative only since numerous modifications and variations therein will be apparent to those skilled in the art.

25

## EXAMPLES

Taking advantage of high avidity resulting from the herein demonstrated specific multivalent interaction between receptor target and recombinant bacteriophage, the

30 examples of the present invention show that a phage clone (Mag-4.1) interacting with the NH<sub>2</sub>-terminal domain of NMDAR1 receptor can be used to monitor the receptor

expression. In the assays below, Mag-4.1 phage was used as a detecting reagent comparable to primary antibodies in immunoassays. As demonstrated in the assays described below, receptor protein can be detected on solid substrates, *e.g.*, on nitrocellulose filters or on transfected mammalian cells. These assays offer opportunities to monitor receptor expression and access binding property profiles of selected peptides prior to more involved affinity maturation.

**Isolation of the specific peptide ligand for NMDA receptor.** To obtain the purified "target" protein for isolating specific peptide ligands, the putative amino-terminal domain of NR1 (N-NR1, a.a 1-561) was cloned and stably expressed in Chinese Hamster Ovary (CHO) cells as a fusion protein, in which the carboxyl-terminal portion contains an epitope tag for monoclonal antibody (mAb8610) (Whitehorn et al., 1995). Prior to panning, the N-NR1 protein was treated with endo-F to remove the glycosylation and the resultant protein preparation was used as a target protein. A phage library with nine random amino peptides was constructed by inserting random oligonucleotides in frame and amino-terminal to the pVIII gene of bacteriophage fl which encodes the major coat protein.

Using N-NR1 as a target protein, Mag-4 phage clone was identified from the random peptide library. After five consecutive rounds of selection and amplification, clones that specifically bind N-NR1 fusion protein were identified by testing the binding of individual phage clone to the immobilized N-NR1 (phage ELISA) (Barrett et al., 1992). Two sequence related phage clones were identified with the deduced sequences of LNDWFITYI (Mag-4.1) (SEQ ID NO:2) and LEAWFLQYI (Mag-4.2) (SEQ ID NO:3). To further test the specificity of phage ELISA experiments, the phage clone of Mag-4.1 was selected and used to test binding to both NH2-terminal domain of NR2A (N-NR2A, a.a. 1-527), an accessory subunit that coassembles with NR1 in vivo (Nakanishi, 1992), and NH2-terminal domain of GluR1 (N-GluR1, a.a. 1-431), a non-NMDA receptor with amino acid sequence homology to NR1 (Hollmann et al., 1989). Figure 1B shows that Mag-4.1 phage binds to N-NR1, but it fails to bind NR2A or N-GluR1. This binding is specific since in the presence of 500 mM of synthetic

Mag-4.1 peptide the binding of Mag-4.1 to N-NR1 was completely inhibited (Figure 1C).

**Detection of N-NR protein by phage-overlay.** Monovalent Mag-4.1 peptide has low affinity (Figure 1C). However, under the conditions of our experiments there are more than 100 copies of Mag-4.1 peptide fused with pVIII coat protein per phage particle. Thus phage particle could have high apparent affinity (avidity) due to multivalent interaction between peptide fusion protein and N-NR1. To test whether this avidity of phage-receptor interaction is sufficient for mediating specific detection, we performed "phage-overlay" assay similar to protein overlay procedure (Li et al., 1992). This binding analysis involves immobilization of target protein on nitrocellulose filter membrane with after separation by SDS-polyacrylamide gel electrophoresis. Figure 2A shows a 5:1 mixture N-NR2A and N-NR1 fusion protein was separated by SDS-PAGE, transferred onto a nitrocellulose filter, and detected by immunoblot using the monoclonal antibody 8610 (mAb8610) (Figure 2, left panel). To detect NR1 protein by Mag-4.1 phage, the immobilized proteins nitrocellulose filter were subjected to denaturation -renaturation procedure using guanidine-HCl (Li et al., 1992). The resultant filter was incubated with Mag-4.1 phage and phage binding were visualized using anti-phage antibody. Figure 2 (right panel) shows that Mag-4.1 phage specifically recognizes N-NR1 protein but not N-NR2A. In addition, the specific binding can be detected by using as little as 20 ng (about 10<sup>9</sup> tu/ml (equivalent 10-12 M)) of N-NR1 protein per lane (Figure 2, right panel, lane 2).

**Detection of N-NR1 protein by phage-immunostain.** Specific detection of N-NR1 by Mag-4.1 phage (Figure 1 and 2) suggests that phage may also be used in way similar to a primary antibody in other immunoassays. We possess two stable Chinese hamster ovary (CHO) cell lines that express N-NR1 on cell surface via GPI anchor (Caras and Weddell, 1989; Whitehorn et al., 1995). These two cell lines have estimated 10<sup>4</sup> (low) and 10<sup>6</sup> (high) copies of N-NR1 per cell. To test whether phage particle coated with specific peptide ligands can be used for immunohistochemistry detection, we incubated Mag-4.1 phage with a 1:1 mixture of the two CHO N-NR1 cell

lines in the presence (Figure 3A) or absence (Figure 3B) of 500 mM Mag-4.1 synthetic peptide. The binding phage particles to cell surface was visualized by anti-phage antibody followed by chromogenic detection (described herein). Figure 3A shows the phage staining on cells expressing N-NR1. The two distinctive staining intensity  
 5 presumably represents the high and low density of receptor present on the two cell lines. This staining signal was completely inhibited when the phage binding was carried out in the presence of 500 mM Mag-4.1 peptide (Figure 3B), suggesting that stain is specific to N-NR1.

10       **Detection of N-NR1 protein by phage-FACS.** To test whether the phage-receptor interaction can be used for fluorescence staining, it is important to access the signal to noise ratio. A double staining experiment was carried out using a cell mixture containing both wildtype CHO cells and CHO cells stably expressing N-NR1. This mixture of cells was stained by mAb8610 (Texas red) and phage  
 15 (fluorescein, green). Figure 4 shows a single view of CHO cells visualized by Nomaski (Figure 4A), mAb8610-Texas red (Figure 4B), and phage-fluorecein (Figure 4C). Indeed, both mAb8610 and Mag-4.1 phage stained the same subset of cells which express N-NR1.

20       To obtain a more quantitative accessment of phage stain, fluorescence activated cell sorting (FACS) was used to detect the phage binding. Figure 5 shows phage staining on four cell preparations: wild type CHO cells (Figure 5A), CHO cells expressing lower N-NR1 receptor per cell (Figure 5B), cells expressing higher N-NR1 receptor per cell (Figure 5C), and a 1:1 mixture of them (Figure 5D). Wild type CHO  
 25 cells gave rise to background signal in Mag-4.1 phage stain (Figure 5A). This signal is similar to the signal obtained by staining the cells with secondary antibody alone. When CHO cells with low density of N-NR1 protein were incubated with Mag-4.1 phage, elevated fluorescence signal was detected (Figure 5B). This signal is specific, as it can be completely inhibited by the synthetic Mag-4.1 peptide (Figure 5C). When a 1:1  
 30 mixture of CHO cells expressing low and high N-NR1 receptor was subjected to the same staining procedure, the two populations of cells were clearly separated as they



have different fluorescence intensity (Figure 5D). Similarly, the staining signal is sensitive to Mag-4.1 peptide (Figure 5E). Together, these results show that it is feasible to use the phage-mediated fluorescence stain to detect the expression of cell surface receptor.

5

- Detection of intact NMDAR in cell culture.** N-NR1 represents a truncated form of NR1, an essential subunit of a pentameric NMDA receptor channel in the brain. To directly test whether Mag-4.1 phage could directly interact with an intact receptor complex, we phage-stained on HEK cells that transiently express NR1 and NR2A
- 10 NMDA receptor subunits. Heteromeric assembly of NR1 and NR2A in HEK cells produces functional NMDA receptor channels with kinetic and pharmacological properties similar to what we have seen in native preparation. When experiments identical to Figure 3 were performed on this preparation, specific Mag-4.1 phage binding signals were observed (Figure 5A and 5B). This staining is specific since only
- 15 a sub-population of cells were stained. The ratio of stained vs unstained cells is identical to that obtained by  $\beta$ -galactosidase stain. In addition, in the presence of 500 mM of synthetic Mag-4.1 peptide the signal was completely inhibited.

**Additional NMDA receptor (NR1) ligands (Mag peptides) isolated.** The following Mag peptides were isolated, in addition to the Mag-4 peptides, by the herein described procedure for isolating specific peptide ligands, and sequence deduced:

<u>Peptide</u>	<u>Amino acid sequence</u>	<u>SEQ ID NO</u>
Mag-2	WDPVFRGCM	5
Mag-3	LGSGGVQRVH	6
Mag-5	MTRATPGMVF	7
Mag-6	TNLRGFWIL	8
Mag-7	FSMDAWFLQYL	9

Native NMDA receptors are likely to be pentameric complexes consisting of NR1 and other subunits such as NR2A. Using the putative agonist binding domain of

NR1 subunit, novel peptides with modulatory activity on native NMDA receptor have been identified. Despite the purified N-NR1 receptor not being in pentamer form, it is clear that some of important structural features of NR1 are retained in the fusion protein. Additional rounds of binding using the reported approach coupled with native  
 5 receptor can be performed to result in peptides with potentially even higher modulatory activity.

### Methods for labeling phage

10 Bacteriophage can be modified in one or a combination of following ways to improve the detection:

(1) phage coat protein (pIII or pVIII) can be engineered to carry recognition sites to permit direct phosphorylation, biotinylation, chemical linkages, etc. to produce phage for convenient detection.

15 (2) phage coat protein (pIII or pVIII) can be linked to known proteins with special properties, e.g., fluorescence protein gene (e.g., green fluorescence protein) and to form a fluorescence phage variant. Such modifications would allow direct detection and enhance phage purification.

(3) phage can be prepared in medium containing radioactive precursors. Thus,  
 20 the resultant phage will be radioactive.

In addition, one can engineer the phage coat to protein with specific properties which reduce the non-specific interactions between phage coat proteins and cells. Such phage variants should be a preferable generic vehicle for all phage staining experiments.

25

### Protocols

**Expression of the NH<sub>2</sub>-terminal domains of NMDA receptor.** Coding sequence for N-NR1 (a.a. 1-561), N-NR2A (a.a. 1-527), N-NR1.spl (a.a. 1- 584), and N-GluR1 (a.a.  
 30 1-431) were obtained by PCR amplification , cloned into eukaryotic cell expression vector driven by the SRa promoter, and confirmed by DNA sequencing. It was then

fused in frame to a nucleotide fragment coding for "AAACLEPYTACDLA  
 PPAGTTDAAHPGRSVVPALL PLLAGTLLLLLETATAP" (SEQ ID NO:1), where the  
 underlined fragment representing the tag for the monoclonal antibody 8610 (mAb8610)  
 and remaining portion is a GPI anchor signal sequence (Caras and Weddell, 1989;  
 5 Whitehorn et al., 1995). The stable transfectants were identified after 2-4 weeks of  
 G418 selection (1 mg/ml). Clones with higher expression were isolated based on the  
 fluorescence intensity and isolated using a FACS-STAR plus (Becton Dickinson).  
 Immunohistochemical detection was carried out using mAb8610 (0.5 mg/ml) followed  
 by 15 mg/ml of secondary antibody (Vector Labs). Imaging was performed using DAB  
 10 based chromagen according to the protocol provided by the manufacturer.

**Peptide Library Construction.** The filamentous phage pVIII gene was chemically  
 synthesized using four overlapping oligonucleotides. This was inserted into the NheI  
 and HindIII sites of the phagemid vector pBAD, which places the expression of the  
 15 pVIII gene under the control of the araB promoter. A cloning site consisting of two  
 non-complementary BstXI sites was included at the 5' end of gene VIII, and this site  
 was used to clone collections of degenerate oligonucleotides for random nine residues  
 (X9) using the half-site strategy as previously described (Cwirla et al., 1990). The  
 resultant library has a complexity of  $1.2 \times 10^{10}$ .

20  
**Phage selection.** Six wells of a 96-well plate were precoated with mAb8610 (5 mg/well)  
 at 37°C for one hour, blocked with 1% BSA for 30 minutes at 37°C. The N-NR1  
 fusion protein was first released by PIPLC digest after extensive washes of cells that  
 express N-NR1. The released N-NR1 protein in the MEM medium was first treated by  
 25 endo-F (1 unit per 10 mg protein) for overnight at 37°C. The treated preparation was  
 affinity-purified by mAb8610 capture on the plates (1 mg of N-NR1 per well) at 22 °C  
 for one hour. The phage binding was carried out by incubating with a total of  $10^{12}$  tu  
 (transducing units) of a X9 library with a complexity of  $1.2 \times 10^{10}$  recombinants. The  
 binding buffer contains 10 mM Tris-HCl (pH 8.1), 100 mM NaCl and 1 mM EDTA for  
 30 3 hours at 4°C. Unbound material was removed with three washes of the binding  
 buffer. The bound phagemid particles were recovered by adding 100 µl of 0.1 M

glycine-HCl (pH 2.2). After 10 minutes elution at room temperature, the elutes were immediately neutralized by mixing with equal volume of 0.1 M Trisbase (pH 9.2). The recovered phagemids were amplified in bacteria, and resultant phage preparations were used for the following round of purification.

5

After five consecutive rounds of selection, the ELISA tests were performed using the supernatant of overnight cultures in the presence of 100 mM Hepes-NaOH (pH7.5) and 100 mM NaCl. The wells were coated with receptors according to the procedure described above. The bound phage particles were first detected by using  
 10 rabbit anti-phage antibody (1: 10,000) followed by alkaline phosphatase (AP)-conjugated secondary antibody (1: 5,000, Sigma). After washing the wells with the binding buffer, 100 ml of developing solution containing 1 mg/ml pNPP (Sigma, MO) in 10 mM diethanolamine-HCl (pH 9.7) was added. The alkaline phosphatase activity was monitored at the wavelength of 408 nm using a microtiter plate reader.

15

**Phage purification.** The bacteria in culture medium containing phage was removed by centrifugation at 5,000 x g for 30 minutes. To precipitate the phage in the resultant supernatant, 0.2 volumes of 20% polyethyleneglycol 8000/2.5 M NaCl were added. After one hour incubation on ice, the precipitated phage was collected by centrifugation  
 20 at 12,000 x g for 30 minutes. The phage pellet was resuspended in 1/10 of starting culture volume and dialyzed overnight against PBS using dialysis membrane with 13 kDa cut-off. The resultant phage preparation was titered.

**Immunoblot.** Proteins were separated by SDS-PAGE and transferred onto a  
 25 nitrocellulose membrane. The nonspecific binding to the membrane was blocked by PBS solution supplemented with 0.1% Tween20, 2% dehydrated milk, and 3% normal goat serum (ICN) for 30 minutes at room temperature. The primary antibodies used were the affinity purified monoclonal antibody 12CA5 (1:5000). The unbound material was removed by three washes with PBS solution supplemented with 0.1% Tween20.  
 30 The binding of primary antibody was detected by adding corresponding peroxidase(HRP)-conjugated secondary antibody (1:5,000) followed by Enhanced

ChemiLuminescence (ECL, Amersham) detection.

**Phage-overlay detection.** Protein lysates containing the NH<sub>2</sub>-terminal domains of glutamate receptors were fractionated by SDS-PAGE and transferred onto a nitrocellulose filter. The following steps were performed at 4 C. The filter containing immobilized proteins was first rinsed in Buffer A (10 mM Hepes-KOH, pH 7.5; 60 mM KCl; 1 mM EDTA, 1 mM 2-mercaptoethanol) for five minutes. It was then exposed for 10 minutes each to Buffer A supplemented with 6M guanidine-HCl at following concentrations: 6M, 3M, 1.5M, 0.75M, 0.38M, 0.19M, 0.1M, 0M. The filter was then incubated for 60 minutes in Buffer A supplemented with 5% of carnation dry milk and 0.05% NP-40. After this incubation, the filter was transferred to the same buffer except 1% carnation added. The phage binding reaction was initiated by first adding the same buffer containing Mag-1 phage with a titer 10<sup>10</sup> tu/ml. After a 12 hour incubation, the unbound phage was removed by three 15 minute wash in Buffer A. The bound phage was detected by adding rabbit anti-phage antibody (1:5,000 dilution) in the same buffer supplemented with 1% normal goat serum and incubating for 1 hour. After three washes to remove unbound immunoglobulin, the HRP-conjugated goat anti-rabbit (Sigma, 1:5,000 dilution) was added to amplify the signals. Finally, the phage binding was detected by ECL using a procedure identical to that described in the section of Immunoblot.

**Immunostain and Fluorescence Activated Cell Sorting (FACS).** The cells were grown on cover glass coated with 1% gelatin. Prior to addition of immunoglobulin, the cells were first washed three times with cold PBS and fixed with 2% paraformaldehyde (PFA) overnight. After washing the cells with cold PBS, the fixed cells were incubated with mAb8610 (1 mg /ml) at room temperature for 1 hour. The binding of mAb8610 was detected by incubating with anti-mouse conjugated with Texas-red (15 mg per ml) 1 hour at 4C. The immunostain was visualized by a Nikon Optiphot-2 fluorescent microscope and images were collected/digitalized using a computer. Cytometry of stained cells were analyzed on either FACScan or FACStar Plus Sorter (Becton-Dickinson, San Jose, CA). An Argon laser is used and tuned at 488 nm. The

green fluorescence was recorded in the FL1 emission channel.

**Phage-immunostain and Phage-FACS detection.** The purified phage preparation was used to stain live cells. For immunostain the cells were washed three times with cold PBS. Phage binding were initiated by incubating with the cells with purified phage (10<sup>10</sup> tu per ml) on ice for 1 hour. After incubation, the unbound phage was removed by three washes of cold PBS supplemented with 1 % bovine serum albumin (BSA). the fixation was carried out using 2% PFA overnight at 4C. The phage-binding was detected first by rabbit anti-phage antibody (1:10,000 dilution) followed by Fluorescein labeled anti-rabbit antibody (10 mg/ml, Vector Lab, CA). The fluorescence labeled cells were visualized under a fluorescence microscope. For Phage-FACS, the cell preparation procedures were identical except the cells were first dislodged from the dishes by incubating with PBS supplemented with 5 mM EDTA.

15

## References

1. Choi, D.W. 1990. Possible mechanisms limiting N-methyl-D-aspartate receptor overactivation and the therapeutic efficacy of N-methyl-D-aspartate antagonists. [Review]. *Stroke*. 21:11120-2
2. Bliss, T.V. & Collingridge, G.L. 1993. A synaptic model of memory: long-term potentiation in the hippocampus. [Review] *Nature* 361, 31-9 (1993).
3. Hollmann, M. & Heinemann, S. 1994. Cloned glutamate receptors. [Review] *Annu Rev Neurosci* 17, 31-108 (1994).
4. Monaghan, D.T., Bridges, R.J. & Cotman, C.W. 1989. The excitatory amino acid receptors: their classes, pharmacology, and distinct properties in the function of the central nervous system. [Review]. *Annu Rev Pharmacol Toxicol* 29, 365-402 (1989).

5. Nakanishi, S. 1992. Molecular diversity of glutamate receptors and implications for brain function. *Science* 258, 597-603
6. Moriyoshi, K., *et al.* 1991. Molecular cloning and characterization of the rat NMDA receptor *Nature* 354, 31-7
7. Westbrook, G.L. & Mayer, M.L. 1987. Micromolar concentrations of  $Zn^{2+}$  antagonize NMDA and GABA responses of hippocampal neurons. *Nature* 328, 640-3.
8. Johnson, J.W. and Ascher, 1987. Glycine potentiates the NMDA response in cultured mouse brain neurons. *Nature*, 325(6104): 529-31.
9. Mayer, M.L., G.L. Westbrook, and P.B. Guthrie, 1984. Voltage-dependent block by  $Mg^{2+}$  of NMDA responses in spinal cord neurones. *Nature*, 309(5965): 261-3.
10. McGurk, J.F., M.V. Bennett, and R.S. Zukin, 1990. Polyamines potentiate responses of N-methyl-D-aspartate receptors expressed in xenopus oocytes. *Proc Natl Acad Sci U S A*, 87(24): 9971-4.
11. Nowak, L., *et al.*, 1984. Magnesium gates glutamate-activated channels in mouse central neurones. *Nature*, 307(5950): 462-5.
12. Peters, S., J. Koh, and D.W. Choi, 1987. Zinc selectively blocks the action of N-methyl-D-aspartate on cortical neurons. *Science*, 236(4801): 589-93.
13. Westbrook, G.L. and M.L. Mayer, 1987. Micromolar concentrations of  $Zn^{2+}$  antagonize NMDA and GABA responses of hippocampal neurons. *Nature*, 328(6131): 640-3.

14. Wong, E.H., et al., 1986. The anticonvulsant MK-801 is a potent N-methyl-D-aspartate antagonist. *Proc Natl Acad Sci U S A*, **83**(18): 7104-8.
15. Lucas, D.R. and J. Newhouse, 1957. The Toxic effect of sodium L-glutamate on the inner layers of retina. *Arch. Ophthalmol.*, **58**: 193-204.
16. Choi, D.W. and S.M. Rothman. 1990. The role of glutamate neurotoxicity in hypoxic-ischemic neuronal death. [Review]. *Annu Rev Neurosci*, **13**: 171-82.
17. Miller, R.J., S.N. Murphy, and S.R. Glaum, 1989. Neuronal  $Ca^{2+}$  channels and their regulation by excitatory amino acids. [Review]. *Ann NY Acad Sci*, **568**: 149-58.
18. Young, A.B., et al., 1983. Glutamate or aspartate as a possible neurotransmitter of cerebral corticofugal fibers in the monkey. *Neurology*, **33**(11): 1513-6.
19. Rothstein, J.D., et al., 1994. Localization of neuronal and glial glutamate transporters. *Neuron*, **13**(3): 713-25.
20. Rothstein, J.D., et al., 1991. Excitatory amino acids in amyotrophic lateral sclerosis: an update [letter]. *Ann Neurol*, **30**(2): 224-5.
21. Rothstein, J.D., L.J. Martin, and R.W. Kuncel, 1992. Decreased glutamate transport by the brain and spinal cord in amyotrophic lateral sclerosis [see comments]. *N Engl J Med*, **326**(22): 1464-8.
22. Rothstein, J.D., et al., 1993. Chronic inhibition of glutamate uptake produces a model of slow neurotoxicity. *Proc Natl Acad Sci U S A*, **90**(14): 6591-5.



23. Storm-Mathisen, J. and O. Ottersen, 1990. Immunocytochemistry of glutamate at the synaptic level. *J Histochem Cytochem*, **38**(12): 1733-43.
24. Storm-Mathisen, J. and O. Ottersen, 1987. Tracing of neurons with glutamate or gamma-aminobutyrate as putative transmitters. [Review]. *Biochem Soc Trans*, **15**(2): 210-3.
25. O'Brien, R.J. and G.D. Fischbach, 1986. Modulation of embryonic chick motoneuron glutamate sensitivity by interneurons and agonists. *J Neurosci*, **6**(11): 3290-6.
26. Tsai, G.C., et al., 1991. Reductions in acidic amino acids and N-acetylaspartylglutamate in amyotrophic lateral sclerosis CNS. *Brain Res*, **556**(1): 151-6.
27. Shaw, P.J., 1994. Excitotoxicity and motor neurone diseases: a review of the evidence. *J. Neurological Sci.*, **124**: 6-13.
28. Johnson, J.W. and Ascher, 1987. Glycine potentiates the NMDA response in cultured mouse brain neurons. *Nature*, **325**(6104): 529-31.
29. Smith, G.P., 1985. Filamentous fusion phage: novel expression vectors that display cloned antigens on the virion surface. *Science*, **228**(4705): 1315-7.
30. Scott, J.K. and G. Smith, 1990. Searching for peptide ligands with an epitope library. *Science*, **249**(4967): 386-90.
31. Cwirla, S.E., et al., 1990. Peptides on phage: a vast library of peptides for identifying ligands. *Proc Natl Acad Sci U S A*, 1990. **87**(16): 6378-82.

32. Devlin, J.J., L.C. Panganiban, and P.E. Devlin. 1990. Random peptide libraries: a source of specific protein binding molecules. *Science*, 249(4967): 404-6.
33. Caras, I.W. & Weddell, G.N. 1989. Signal peptide for protein secretion directing glycopospholipid membrane anchor attachment. *Science* 243, 1196-8 (1989).
34. Chazot, P.L., Cik, M. & Stephenson, F.A. 1992. Immunological detection of the NMDAR1 glutamate receptor subunit expressed in embryonic kidney 293 cells and in rat brain. *J Neurochem* 59, 1176-8.
35. Barrett, R.W., *et al.*, 1992. Selective enrichment and characterization of high affinity ligands from collections of random peptides on filamentous phage. *Anal Biochem*, 204(2): p. 357-64.
36. Haack, J.A., *et al.*, 1990. Conantokin-T. A gamma-carboxyglutamate containing peptide with N-methyl-d-aspartate antagonist activity. *J Biol Chem*, 265(11): 6025-9.
37. Huettner, J.E. and B.P. Bean, 1988. Block of N-methyl-D-aspartate-activated current by the anticonvulsant MK-801: selective binding to open channels. *Proc Natl Acad Sci U S A*, 85(4): p. 1307-11.
38. Mori, H., *et al.*, 1992. Identification by mutagenesis of a Mg(2+)-block site of the NMDA receptor channel. *Nature*, 358(6388): 673-5.
39. Monyer, H., *et al.*, 1992 Heteromeric NMDA receptors: molecular and functional distinction of subtypes. *Science*, 256(5060): 1217-21.
40. Kutsuwada, T., *et al.*, 1992. Molecular diversity of the NMDA receptor channel [see comments]. *Nature*, 358(6381): 36-41.

41. Tingley, W.G., *et al.*, 1993 Regulation of NMDA receptor phosphorylation by alternative splicing of the C-terminal domain. *Nature*, 1993. 364(6432): 70-3.
42. Cik, M., P.L. Chazot, and F.A. Stephenson. 1993. Optimal expression of cloned NMDAR1/NMDAR2A heteromeric glutamate receptors: a biochemical characterization. *Biochem J*, 1993. 296(Pt 3): 877-83.
43. Li, M., Jan, Y.N. & Jan, L.Y. 1992. Specification of subunit assembly by the hydrophilic amino-terminal domain of the Shaker potassium channel. *Science* 257, 1225-30.
44. Blond-Elguindi, S., Cwirla, S. E., Dower, W. J., Lipshutz, R. J., Sprang, S. R., Sambrook, J. F., and Gething, M. J. (1993). Affinity panning of a library of peptides displayed on bacteriophages reveals the binding specificity of BiP. *Cell* 75, 717-28.
45. Dedman, J. R., Kaetzel, M. A., Chan, H. C., Nelson, D. J., and Jamieson, G., Jr. (1993). Selection of targeted biological modifiers from a bacteriophage library of random peptides. The identification of novel calmodulin regulatory peptides. *J Biol Chem* 268, 23025-30.
46. Greenamyre, J. T., Penney, J. B., Young, A. B., D'amato, C. J., Hicks, S. P., and Shoulson, I. (1985). Alterations in L-glutamate binding in Alzheimer's and Huntington's diseases. *Science* 227.
47. Hollmann, M., O'Shea-Greenfield, A., Rogers, S. W., and Heinemann, S. (1989). Cloning by functional expression of a member of the glutamate receptor family. *Nature* 342, 643-8.
48. Kay, B. K., Adey, N. B., He, Y. S., Manfredi, J. P., Mataragnon, A. H., and Fowlkes, D. M. (1993). An M13 phage library displaying random 38-amino-acid

peptides as a source of novel sequences with affinity to selected targets. *Gene* 128, 59-65.

49. Rice, G. C., Goeddel, D. V., Cachianes, G., Woronicz, J., Chen, E. Y., Williams, S. R., and Leung, D. W. (1992). Random PCR mutagenesis screening of secreted proteins by direct expression in mammalian cells. *Proc. Natl. Acad. Sci. USA* 89, 5467-5471.

50. Sparks, A. B., Quilliam, L. A., Thorn, J. M., Der, C. J., and Kay, B. K. (1994). Identification and characterization of Src SH3 ligands from phage-displayed random peptide libraries. *J Biol Chem* 269, 23853-6.

51. Whitehorn, E. A., Tate, E., Yanofsky, S. D., Kochersperger, L., Davis, A., Mortensen, R. B., Yonkovich, S., Bell, K., Dower, W. J., and Barrett, R. W. (1995). A generic method for expression and use of "tagged" soluble versions of cell surface receptors. *Biotechnology* 13, 1215-1219.

Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

Although the present process has been described with reference to specific details of certain embodiments thereof, it is not intended that such details should be regarded as limitations upon the scope of the invention except as and to the extent that they are included in the accompanying claims.

(1) GENERAL INFORMATION:

- (2) INFORMATION FOR SEQ ID NO:1:

- (ii) MOLECULE TYPE: peptide

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Ala Ala Ala Cys Leu Glu Pro Tyr Thr Ala Cys Asp Leu Ala Pro Pro  
1 5 10 15

Ala Gly Thr Thr Asp Ala Ala His Pro Gly Arg Ser Val Val Pro Ala  
20 25 30

Leu Leu Pro Leu Leu Ala Gly Thr Leu Leu Leu Leu Glu Thr Ala Thr  
           35                                  40                                  45

Ala Pro  
       50

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:  
       (A) LENGTH: 9 amino acids  
       (B) TYPE: amino acid  
       (C) STRANDEDNESS: not relevant  
       (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Leu Asn Asp Trp Phe Ile Thr Tyr Ile  
   1                                  5

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:  
       (A) LENGTH: 9 amino acids  
       (B) TYPE: amino acid  
       (C) STRANDEDNESS: not relevant  
       (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Leu Glu Ala Trp Phe Leu Gln Tyr Ile  
   1                                  5

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:  
       (A) LENGTH: 15 amino acids  
       (B) TYPE: amino acid  
       (C) STRANDEDNESS: not relevant  
       (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Gly Gly Gly Leu Asn Asp Trp Phe Ile Thr Tyr Ile Gly Gly Gly  
   1                                  5                                  10                                  15

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Trp Asp Pro Ala Val Phe Arg Gly Cys Met  
1                      5                      10

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 10 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: not relevant
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Leu Gly Ser Gly Gly Val Gln Arg Val His  
1                      5                      10

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 10 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: not relevant
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met Thr Arg Ala Thr Pro Gly Met Val Phe  
1                      5                      10

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 11 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: not relevant
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Thr Asn Leu Arg Arg Gly Phe Trp Ile Ile Leu  
1 5 10

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Phe Ser Met Asp Ala Trp Phe Leu Gln Tyr Leu  
1 5 10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2